

Time-series Spectroscopy of Pulsating sdB Stars: PG1605+072¹

S. J. O'Toole and T. R. Bedding

School of Physics, University of Sydney, NSW 2006, Australia

and

H. Kjeldsen

Theoretical Astrophysics Center, Aarhus University, DK-8000, Aarhus C, Denmark

and

T. C. Teixeira

The MONS Project Team, Institute of Physics and Astronomy, Aarhus University,
DK-8000 Aarhus C, Denmark

and

G. Roberts, F. van Wyk and D. Kilkeny

South African Astronomical Observatory, PO Box 9, Observatory 7935, South Africa

and

N. D'Cruz and I. K. Baldry

School of Physics, University of Sydney, NSW 2006, Australia

Received _____; accepted _____

ABSTRACT

We report the detection of velocity variations in the pulsating sdB star, PG 1605+072. Oscillations are detected at the same frequencies found from photometry and have amplitudes of up to 14 km s^{-1} for $\text{H}\beta$. The strongest oscillation found in previous photometric observations is not evident in our spectroscopy or photometry, and may be absent due to beating of closely spaced modes. Phase differences between spectroscopy and B magnitude photometry imply that maximum brightness occurs not long after maximum radius. We have also found evidence of variation in the observed amplitudes of five Balmer lines, with a decrease in amplitude of the strongest mode blueward from $\text{H}\beta$. This effect is not expected and a longer time-series will be needed to clarify it.

Subject headings: stars: interiors — stars: oscillations — subdwarfs

1. Introduction

The discovery of a class of pulsating hot subdwarfs (sdBs; Kilkenney et al. 1997) has created an interest in asteroseismology of evolved stars. Subdwarfs are believed to lie on the Extreme Horizontal Branch (Saffer et al. 1994), although their formation and evolution is still uncertain. Pulsations in sdBs were predicted theoretically by Charpinet et al. (1996) just prior to their discovery, and about 20 pulsators have been discovered to date using time-series photometry (see Koen et al. 1999 for a review). The stars (also known

¹Based on observations made with the Danish 1.54 m telescope at ESO, La Silla, Chile, from the South African Astronomical Observatory (SAAO), and from Mt Stromlo Observatory, Australia.

as EC14026 stars, after the prototype) are believed to pulsate in p modes and typically have periods 100–200 s and semi-amplitudes <10 mmag. Models indicate an opacity bump associated with iron ionisation may be the driving mechanism of the pulsations (Charpinet et al. 1996, 1997), although recent studies by Koen et al. (1999) find no well-defined instability strip in the $\log g - T_{\text{eff}}$ plane.

Combining time-series photometry with time-series spectroscopy should help to give a better understanding of the pulsations in these stars. We have shown the feasibility of time-series spectroscopy with observations of PG1605+072 on a 1.5 m telescope over 2 hrs (O’Toole et al. 2000). Jeffery & Pollacco (2000) made similar observations of KPD2109+4401 (5 hrs) and PB8783 (6 hrs) using the 4.2m William Herschel Telescope. In all three cases, velocity variations were detected at the published photometric frequencies. However, sdB stars are multimode pulsators and longer time series are needed to resolve the individual modes and measure their amplitudes. We present here the first extended time-series spectroscopy of a pulsating sdB star.

Our target was PG1605+072 which has the most extreme properties of all the pulsating sdBs detected so far, with high amplitudes (up to ~ 25 mmag) and long periods (~ 500 s). Because of its low gravity ($\log g \sim 5.25$), this star is believed to have evolved off the Horizontal Branch and may oscillate in g , as well as p modes (Kilkenny et al. 1999). The power spectrum is complex, with up to 55 indentifiable frequencies. PG1605+072 displays considerable rotation ($v \sin i = 39 \text{ km s}^{-1}$), possibly leading to unequally spaced multiplet components (Heber et al. 1999). Unlike several of the pulsating sdBs, multicolour photometry gives no indication of a cool companion (Koen et al. 1998).

2. Observations

We obtained medium-resolution spectra of PG1605+072 using the DFOSC spectrograph on the Danish 1.54 m telescope at La Silla, Chile and the coudé spectrograph (A grating) on the 74-inch (1.88 m) telescope at Mt Stromlo, Australia. The observations were made on 7 nights over an 11-day period in July and August, 1999 (see Table 1). To supplement the spectroscopy, we obtained time-series photometry at the Sutherland site of the South African Astronomical Observatory (see Table 2).

EDITOR: PLACE TABLE 1 HERE.

EDITOR: PLACE TABLE 2 HERE.

The La Silla data consisted of single-order spectra projected onto a 2K LORAL CCD - pixel-binning (to reduce R.O.N.) and windowing (to reduce readout time) gave 66 x 500 pixel spectra with a total wavelength range of 3700–5000 Å and a dispersion of 1.65 Å pixel⁻¹. The resolution was 2.8 Å, set by a slit width of 1.5 arcsec. The exposure time was 46 s, with a dead time of about 15 s. The average number of photons per Å in each spectrum was about 2200.

The Mt Stromlo data consisted of single-order spectra projected onto a SITe 2Kx4K CCD, with a wavelength range of 3800–5000 Å. The slit width was ~4 arcsec and the dispersion was 0.60 Å pixels⁻¹, leading to a resolution similar to that of the La Silla data. The exposure time was 50 s with a dead time of around 25 s.

The photometry from Sutherland was taken using the Modular Photometer with a Johnson *B* filter and exposure times of 20 s. The GaAs photomultiplier tube in this photometer has a good red sensitivity, and using the blue filter causes the passband to resemble that of a blue-sensitive photomultiplier tube (O’Donoghue et al. 1998).

3. Reductions

Standard methods for bias subtraction, flat-fielding and background scattered-light subtraction were used. Non-linearities on the CCD image were only important in the Stromlo data, and were corrected immediately after bias subtraction (Baldry et al. 1998, see). One-dimensional spectra were extracted using the Optimal Extraction method (Horne 1986), a variance weighting system for CCD columns. A 3rd order polynomial was fitted to the continuum level in both data sets, and the spectra were normalised to a continuum value of unity.

The spectrum of PG1605+072 is dominated by Balmer lines - see e.g. Figure 9 of Koen et al. (1998). A cross-correlation technique was used to determine the Doppler shift for five Balmer lines ($H\beta$, $H\gamma$, $H\delta$, $H\epsilon$ and $H8$), relative to a template spectrum (the average of 20 high-quality spectra). The spectra and template were prepared in the manner described by Baldry et al. (1998). It should be noted that the time-series for $H\gamma$ does not contain any Mt Stromlo data, because that line was contaminated by a bad CCD column.

Several jumps were found in each time-series, particularly in the La Silla data, occurring every 30-40 observations, corresponding to breaks where the He+Ne calibration spectra were taken. We decided to process each of these groups separately with a template spectrum created for each group. In removing the jumps, we have effectively high-pass filtered the dataset, also removing any instrumental drifts present.

EDITOR: PLACE FIGURE 1 HERE.

E1

^{E1}NOTE TO EDITOR: rawvelocity.ps should be over two columns

The photometric data were reduced by subtracting the sky background, correcting for atmospheric extinction and normalising to the mean intensity of the run, following a method similar to O’Donoghue et al. (1997, 1998). Slow drifts were not corrected for in the photometry.

The $H\beta$ velocity curves and B magnitude light curves for each night of observations are shown in Figure 1. Oscillations are not as evident in our velocity curve as they are in the light curve, since the signal-to-noise per measurement is lower.

As is common with time-series measurements, the quality of the data varies considerably through the data set. We therefore performed frequency analysis using a weighted Fourier Transform, where weights were assigned according to the local rms scatter (Kjeldsen & Frandsen 1992).

4. Results and Discussion

The amplitude spectra of $H\beta$ and $H\gamma$ are shown in the top two panels on the left of Figure 2. The white noise level is 1.07 km s^{-1} for $H\beta$ and 0.93 km s^{-1} for $H\gamma$. Oscillations are clearly visible at $2102 \mu\text{Hz}$ and $2743 \mu\text{Hz}$ ($2742.72 \mu\text{Hz}$ in Kilkenney et al. 1999) in both lines but at $1891 \mu\text{Hz}$ ($1891.42 \mu\text{Hz}$) only in $H\gamma$. It is unclear whether this last effect is due to the low amplitude of the mode in $H\beta$ due to beating between closely spaced modes. The bottom two panels show our B magnitude and white light amplitude spectra (from a 15-day multisite campaign by Kilkenney et al. (1999) in May, 1997; white light observations were obtained using a CuSO_4 filter) respectively, and are included for comparison. The three modes previously mentioned are all present in the B magnitude spectrum. We believe that the highest amplitude peak (14 km s^{-1} for $H\beta$) at $2102 \mu\text{Hz}$ may be a combination of the $2101.65 \mu\text{Hz}$ and $2103.28 \mu\text{Hz}$ frequencies found by Kilkenney et al. – a longer time-series is

needed to clarify this.

EDITOR: PLACE FIGURE 2 HERE.

E2

The panels on the right of Figure 2 show the 2050–2150 μHz region in greater detail. The highest amplitude mode found by Kilkenny et al. at 2075.76 μHz is not present in any of our observations. It is not clear whether this is due to variation in amplitudes over the 2.2 years between observations, or to beating between very closely spaced modes.

The B magnitude and white light amplitudes differ by around 40%, which is to be expected in such hot, blue stars, considering that the effective wavelength of white light is redder than the B band. If oscillation phase is wavelength dependent, then this would also reduce the amplitude in white light (which is a much broader “band” than B).

4.1. Oscillation Amplitudes

In this Letter we present a qualitative comparison of luminosity and velocity amplitudes. All amplitudes were measured by the height of their peak. A more detailed quantitative study will be the subject of a subsequent paper.

Kjeldsen & Bedding (1995) derived a relationship between luminosity amplitude at a given wavelength and velocity amplitude for classical and solar-like oscillations. Using their scaling law, which assumes a black body spectrum and adiabatic oscillations, we find that the velocity amplitudes in PG1605+072 are 2–3 times lower than expected. However a black body and adiabatic assumption is far from valid in such a hot star

^{E2}NOTE TO EDITOR: compspec.ps should be over two columns

when one is considering amplitudes in visual. In a pure adiabatic assumption there is a 180° phase difference between temperature and stellar radius (with maximum brightness coinciding with minimum radius). For the mode at 2.742 mHz (which do not appear to be contaminated by closely spaced modes), we find a phase difference of $(-75 \pm 10)^\circ$ implying that maximum brightness is slightly after maximum radius. More data are needed to clarify this result.

EDITOR: PLACE FIGURE 3 HERE.

E3

There is evidence to suggest that the velocity amplitude is wavelength dependent. Figure 3 shows that the amplitude of oscillations at two frequencies decreases blueward of $H\beta$. The amplitude of H8 is almost 25% smaller than that of $H\beta$. If confirmed, this would be puzzling, given that all the Balmer lines are formed in the same part of the atmosphere.

5. Conclusions

The results presented here show that times-series spectroscopy of sdBs can be done using a medium-size (1.5 m) telescope. We have detected oscillations in velocity at previously published photometric frequencies. A longer times-series would allow most of the closely spaced modes in PG1605+072 to be resolved and thus help to confirm the variation of velocity amplitude with wavelength. Further studies may also determine whether the amplitudes of modes in this star are variable over time, or whether the apparent shift in primary pulsation mode is due to beating. We will also examine the behaviour of equivalent width and other line profile variations.

^{E3}NOTE TO EDITOR: wave-vs-amp.dat.ps should be over one column

This work was supported by an Australian Postgraduate Award (SJOT), the Australian Research Council, the Danish National Science Research Council through its Center for Ground-based Observational Astronomy, and the Danish National Research Foundation through its establishment of the Theoretical Astrophysics Center.

REFERENCES

- Baldry, I. K., Bedding, T. R., Viskum, M., Kjeldsen, H., & Frandsen, S., 1998, MNRAS, 295, 33
- Charpinet, S., Fontaine, G., Brassard, P., & Dorman, B., 1996, ApJ, 471, L103
- Charpinet, S., Fontaine, G., Brassard, P., Chayer, P., Rogers, F. J., Iglesias, C. A., & Dorman, B., 1997, ApJ, 483, L123
- Heber, U., Reid, I. N., & Werner, K., August 1999, A&A, 348, L25
- Horne, K., June 1986, PASP, 98, 609–617
- Jeffery, C. S., & Pollacco, D., 2000, MNRAS, submitted
- Kilkenny, D., Koen, C., O’Donoghue, D., & Stobie, R. S., 1997, MNRAS, 285, 640
- Kilkenny, D., Koen, C., O’Donoghue, D., Van Wyk, F., Larson, K. A., Shobbrook, R., Sullivan, D. J., Burleigh, M. R., Dobbie, P. D., & Kawaler, S. D., 1999, MNRAS, 303, 525
- Kjeldsen, H., & Bedding, T. R., 1995, A&A, 293, 87
- Kjeldsen, H., & Frandsen, S., 1992, PASP, 104, 413
- Koen, C., O’Donoghue, D., Kilkenny, D., Lynas-Gray, A. E., Marang, F., & Van Wyk, F., 1998, MNRAS, 296, 317
- Koen, C., O’Donoghue, D., Kilkenny, D., Stobie, R. S., & Saffer, R. A., 1999, MNRAS, 306, 213
- O’Donoghue, D., Lynas-Gray, A. E., Kilkenny, D., Stobie, R. S., & Koen, C., 1997, MNRAS, 285, 657

- O'Donoghue, D., Koen, C., Lynas-Gray, A. E., Kilkenney, D., & Van Wyk, F., 1998, MNRAS, 296, 306
- O'Toole, S. J., Teixeira, T. C., Bedding, T. R., & Kjeldsen, H., 2000. In: Kurtz, D., & Szabados, L. (eds.), *The Impact of Large Scale Surveys on Pulsating Star Research*, *IAU Colloquium 176*, in press
- Saffer, R. A., Bergeron, P., Koester, D., & Liebert, J., 1994, ApJ, 432, 351

Table 1: Spectroscopic observations of PG1605+072. LS = La Silla; MS = Mt Stromlo.

UT-date	Site	No. of hours	No. of spectra
1999 July 23–4	LS	4.01	210
1999 July 24–5	LS	1.91	105
1999 July 25–6	LS	5.46	280
1999 July 26–7	LS	5.12	280
1999 July 30–1	LS	5.09	260
1999 July 31	MS	3.61	168
1999 July 31–1	LS	5.23	280
1999 Aug 01	MS	3.30	155
1999 Aug 01–2	LS	4.92	280
total		38.65	2018

Table 2: Photometric observations of PG1605+072, taken at the Sutherland site of the SAAO.

UT-date	No. of hours	No. of obs.
1999 July 20	3.93	666
1999 July 21	1.07	182
1999 July 23	4.17	679
1999 July 24	1.27	211
1999 July 28	3.77	597
1999 July 29	1.19	199
1999 July 31	1.24	210
1999 Aug 01	2.80	454
total	19.44	3198




Fig. 1.— Combined raw $H\beta$ velocity and Johnson B light curve for PG1605+072. Velocity points (in km s^{-1}) are large dots while photometry points are small dots. Photometry in this Letter is presented in ‘millimodulation units’ or mmi, where the light curve is normalized by its own mean level (see Kilkenney et al. 1999). Observations were made on JD 2451 380-393 (see Tables 1 and 2). Note that positive velocity is redshifted by definition.




Fig. 2.— Amplitude spectrum of PG1605+072 in the 1000–5000 μHz range. Panels on the left from top to bottom: $\text{H}\beta$ velocity; $\text{H}\gamma$ velocity (with no Stromlo data); B band photometry; Photometry by Kilkenney et al. (1999) in May, 1997 – the dotted lines represent the 10 strongest modes found. Panels on the right show greater detail centred around 2100 μHz .




Fig. 3.— Velocity amplitude vs wavelength for the 5 Balmer lines studied. The group of unresolved modes at 2.10 mHz is represented by diamonds, while the mode at 2.74 mHz is represented by triangles.